

519,596, published Dec. 23, 1992). These "humanized" molecules are designed to minimize unwanted immunological response toward rodent antihuman antibody molecules which limits the duration and effectiveness of therapeutic applications of those moieties in human recipients.

5 As used herein, the terms "veneered FRs" and "recombinantly veneered FRs" refer to the selective replacement of FR residues from, e.g., a rodent heavy or light chain V region, with human FR residues in order to provide a xenogeneic molecule comprising an antigen-binding site which retains substantially all of the native FR polypeptide folding structure. Veneering techniques are based on the understanding that  
10 the ligand binding characteristics of an antigen-binding site are determined primarily by the structure and relative disposition of the heavy and light chain CDR sets within the antigen-binding surface. Davies et al. (1990) *Ann. Rev. Biochem.* 59:439-473. Thus, antigen binding specificity can be preserved in a humanized antibody only wherein the CDR structures, their interaction with each other, and their interaction with the rest of  
15 the V region domains are carefully maintained. By using veneering techniques, exterior (e.g., solvent-accessible) FR residues which are readily encountered by the immune system are selectively replaced with human residues to provide a hybrid molecule that comprises either a weakly immunogenic, or substantially non-immunogenic veneered surface.

20 The process of veneering makes use of the available sequence data for human antibody variable domains compiled by Kabat et al., in *Sequences of Proteins of Immunological Interest*, 4th ed., (U.S. Dept. of Health and Human Services, U.S. Government Printing Office, 1987), updates to the Kabat database, and other accessible U.S. and foreign databases (both nucleic acid and protein). Solvent accessibilities of V  
25 region amino acids can be deduced from the known three-dimensional structure for human and murine antibody fragments. There are two general steps in veneering a murine antigen-binding site. Initially, the FRs of the variable domains of an antibody molecule of interest are compared with corresponding FR sequences of human variable domains obtained from the above-identified sources. The most homologous human V  
30 regions are then compared residue by residue to corresponding murine amino acids. The

residues in the murine FR which differ from the human counterpart are replaced by the residues present in the human moiety using recombinant techniques well known in the art. Residue switching is only carried out with moieties which are at least partially exposed (solvent accessible), and care is exercised in the replacement of amino acid residues which may have a significant effect on the tertiary structure of V region domains, such as proline, glycine and charged amino acids.

In this manner, the resultant "veneered" murine antigen-binding sites are thus designed to retain the murine CDR residues, the residues substantially adjacent to the CDRs, the residues identified as buried or mostly buried (solvent inaccessible), the residues believed to participate in non-covalent (e.g., electrostatic and hydrophobic) contacts between heavy and light chain domains, and the residues from conserved structural regions of the FRs which are believed to influence the "canonical" tertiary structures of the CDR loops. These design criteria are then used to prepare recombinant nucleotide sequences which combine the CDRs of both the heavy and light chain of a murine antigen-binding site into human-appearing FRs that can be used to transfect mammalian cells for the expression of recombinant human antibodies which exhibit the antigen specificity of the murine antibody molecule.

In another embodiment of the invention, monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include  $^{90}\text{Y}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{211}\text{At}$ , and  $^{212}\text{Bi}$ . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, Pseudomonas exotoxin, Shigella toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (e.g., covalently bonded) to a suitable monoclonal antibody either directly or indirectly (e.g., via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulphydryl group, on one may be capable of reacting with a carbonyl-

containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, thiol groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitzer), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be

coupled directly to an antibody molecule, or linkers that provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (e.g., U.S. Patent No. 4,567,234, to Kato et al.), peptides and polysaccharides such as aminodextrans (e.g., U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (e.g., U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiobalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

#### T Cell Compositions

The present invention, in another aspect, provides T cells specific for a tumor polypeptide disclosed herein, or for a variant or derivative thereof. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be isolated from bone marrow, peripheral blood, or a fraction of bone marrow or peripheral blood of a patient, using a commercially available cell separation system, such as the Isoplex™ System, available from Nexell Therapeutics, Inc. (Irvine, CA; see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans, non-human mammals, cell lines or cultures.

T cells may be stimulated with a polypeptide, polynucleotide encoding a polypeptide and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the polypeptide of interest. Preferably, a tumor

polypeptide or polynucleotide of the invention is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for a polypeptide of the present invention if the T cells specifically proliferate, secrete cytokines or kill target cells coated with the polypeptide or expressing a gene encoding the polypeptide. T cell specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994. Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (e.g., by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with a tumor polypeptide (100 ng/ml - 100 µg/ml, preferably 200 ng/ml - 25 µg/ml) for 3 - 7 days will typically result in at least a two fold increase in proliferation of the T cells. Contact as described above for 2-5 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (e.g., TNF or IFN-γ) is indicative of T cell activation (see Coligan et al., *Current Protocols in Immunology*, vol. 1, Wiley Interscience (Greene 1998)). T cells that have been activated in response to a tumor polypeptide, polynucleotide or polypeptide-expressing APC may be CD4<sup>+</sup> and/or CD8<sup>+</sup>. Tumor polypeptide-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient, a related donor or an unrelated donor, and are administered to the patient following stimulation and expansion.

For therapeutic purposes, CD4<sup>+</sup> or CD8<sup>+</sup> T cells that proliferate in response to a tumor polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to a tumor polypeptide, or a short peptide corresponding to an immunogenic portion of such a polypeptide, with or

without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize a tumor polypeptide. Alternatively, one or more T cells that proliferate in the presence of the tumor polypeptide can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting  
5 dilution.

#### Pharmaceutical Compositions

In additional embodiments, the present invention concerns formulation of one or more of the polynucleotide, polypeptide, T-cell and/or antibody compositions disclosed herein in pharmaceutically-acceptable carriers for administration to a cell or  
10 an animal, either alone, or in combination with one or more other modalities of therapy.

It will be understood that, if desired, a composition as disclosed herein may be administered in combination with other agents as well, such as, *e.g.*, other proteins or polypeptides or various pharmaceutically-active agents. In fact, there is virtually no limit to other components that may also be included, given that the  
15 additional agents do not cause a significant adverse effect upon contact with the target cells or host tissues. The compositions may thus be delivered along with various other agents as required in the particular instance. Such compositions may be purified from host cells or other biological sources, or alternatively may be chemically synthesized as described herein. Likewise, such compositions may further comprise substituted or  
20 derivatized RNA or DNA compositions.

Therefore, in another aspect of the present invention, pharmaceutical compositions are provided comprising one or more of the polynucleotide, polypeptide, antibody, and/or T-cell compositions described herein in combination with a physiologically acceptable carrier. In certain preferred embodiments, the  
25 pharmaceutical compositions of the invention comprise immunogenic polynucleotide and/or polypeptide compositions of the invention for use in prophylactic and therapeutic vaccine applications. Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Generally, such compositions will comprise one or more

polynucleotide and/or polypeptide compositions of the present invention in combination with one or more immunostimulants.

It will be apparent that any of the pharmaceutical compositions described herein can contain pharmaceutically acceptable salts of the polynucleotides and polypeptides of the invention. Such salts can be prepared, for example, from pharmaceutically acceptable non-toxic bases, including organic bases (e.g., salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (e.g., sodium, potassium, lithium, ammonium, calcium and magnesium salts).

In another embodiment, illustrative immunogenic compositions, e.g., vaccine compositions, of the present invention comprise DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the polynucleotide may be administered within any of a variety of delivery systems known to those of ordinary skill in the art. Indeed, numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198, 1998, and references cited therein. Appropriate polynucleotide expression systems will, of course, contain the necessary regulatory DNA regulatory sequences for expression in a patient (such as a suitable promoter and terminating signal). Alternatively, bacterial delivery systems may involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface or secretes such an epitope.

Therefore, in certain embodiments, polynucleotides encoding immunogenic polypeptides described herein are introduced into suitable mammalian host cells for expression using any of a number of known viral-based systems. In one illustrative embodiment, retroviruses provide a convenient and effective platform for gene delivery systems. A selected nucleotide sequence encoding a polypeptide of the present invention can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to a subject. A number of illustrative retroviral systems have been described (e.g., U.S. Pat. No. 5,219,740; Miller and Roman (1989) *BioTechniques* 7:980-990; Miller, A. D. (1990) *Human Gene Therapy* 1:5-14; Scarpa et al. (1991) *Virology* 180:849-852; Burns

et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:8033-8037; and Boris-Lawrie and Temin (1993) *Curr. Opin. Genet. Develop.* 3:102-109.

In addition, a number of illustrative adenovirus-based systems have also been described. Unlike retroviruses which integrate into the host genome, adenoviruses  
5 persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis (Haj-Ahmad and Graham (1986) *J. Virol.* 57:267-274; Bett et al. (1993) *J. Virol.* 67:5911-5921; Mittereder et al. (1994) *Human Gene Therapy* 5:717-729; Seth et al. (1994) *J. Virol.* 68:933-940; Barr et al. (1994) *Gene Therapy* 1:51-58; Berkner, K. L. (1988) *BioTechniques* 6:616-629; and Rich et al. (1993) *Human Gene Therapy* 4:461-  
10 476).

Various adeno-associated virus (AAV) vector systems have also been developed for polynucleotide delivery. AAV vectors can be readily constructed using techniques well known in the art. See, e.g., U.S. Pat. Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski et al.  
15 (1988) *Molec. Cell. Biol.* 8:3988-3996; Vincent et al. (1990) *Vaccines* 90 (Cold Spring Harbor Laboratory Press); Carter, B. J. (1992) *Current Opinion in Biotechnology* 3:533-539; Muzyczka, N. (1992) *Current Topics in Microbiol. and Immunol.* 158:97-129; Kotin, R. M. (1994) *Human Gene Therapy* 5:793-801; Shelling and Smith (1994) *Gene Therapy* 1:165-169; and Zhou et al. (1994) *J. Exp. Med.* 179:1867-1875.

20 Additional viral vectors useful for delivering the polynucleotides encoding polypeptides of the present invention by gene transfer include those derived from the pox family of viruses, such as vaccinia virus and avian poxvirus. By way of example, vaccinia virus recombinants expressing the novel molecules can be constructed as follows. The DNA encoding a polypeptide is first inserted into an  
25 appropriate vector so that it is adjacent to a vaccinia promoter and flanking vaccinia DNA sequences, such as the sequence encoding thymidine kinase (TK). This vector is then used to transfect cells which are simultaneously infected with vaccinia. Homologous recombination serves to insert the vaccinia promoter plus the gene encoding the polypeptide of interest into the viral genome. The resulting TK<sub>sup</sub>(-)



recombinant can be selected by culturing the cells in the presence of 5-bromodeoxyuridine and picking viral plaques resistant thereto.

A vaccinia-based infection/transfection system can be conveniently used to provide for inducible, transient expression or coexpression of one or more polypeptides described herein in host cells of an organism. In this particular system, cells are first infected *in vitro* with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the polynucleotide or polynucleotides of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into polypeptide by the host translational machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation products. See, *e.g.*, Elroy-Stein and Moss, *Proc. Natl. Acad. Sci. USA* (1990) 87:6743-6747; Fuerst *et al.* *Proc. Natl. Acad. Sci. USA* (1986) 83:8122-8126.

Alternatively, avipoxviruses, such as the fowlpox and canarypox viruses, can also be used to deliver the coding sequences of interest. Recombinant avipox viruses, expressing immunogens from mammalian pathogens, are known to confer protective immunity when administered to non-avian species. The use of an Avipox vector is particularly desirable in human and other mammalian species since members of the Avipox genus can only productively replicate in susceptible avian species and therefore are not infective in mammalian cells. Methods for producing recombinant Avipoxviruses are known in the art and employ genetic recombination, as described above with respect to the production of vaccinia viruses. See, *e.g.*, WO 91/12882; WO 89/03429; and WO 92/03545.

Any of a number of alphavirus vectors can also be used for delivery of polynucleotide compositions of the present invention, such as those vectors described in U.S. Patent Nos. 5,843,723; 6,015,686; 6,008,035 and 6,015,694. Certain vectors based on Venezuelan Equine Encephalitis (VEE) can also be used, illustrative examples of which can be found in U.S. Patent Nos. 5,505,947 and 5,643,576.

Moreover, molecular conjugate vectors, such as the adenovirus chimeric vectors described in Michael et al. *J. Biol. Chem.* (1993) 268:6866-6869 and Wagner et al. *Proc. Natl. Acad. Sci. USA* (1992) 89:6099-6103, can also be used for gene delivery under the invention.

- 5 Additional illustrative information on these and other known viral-based delivery systems can be found, for example, in Fisher-Hoch et al., *Proc. Natl. Acad. Sci. USA* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242;  
10 WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfield et al., *Science* 252:431-434, 1991; Kolls et al., *Proc. Natl. Acad. Sci. USA* 91:215-219, 1994; Kass-Bisler et al., *Proc. Natl. Acad. Sci. USA* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993.

- In certain embodiments, a polynucleotide may be integrated into the  
15 genome of a target cell. This integration may be in the specific location and orientation via homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the polynucleotide may be stably maintained in the cell as a separate, episomal segment of DNA. Such polynucleotide segments or "episomes" encode sequences sufficient to  
20 permit maintenance and replication independent of or in synchronization with the host cell cycle. The manner in which the expression construct is delivered to a cell and where in the cell the polynucleotide remains is dependent on the type of expression construct employed.

- In another embodiment of the invention, a polynucleotide is  
25 administered/delivered as "naked" DNA, for example as described in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

- In still another embodiment, a composition of the present invention can  
30 be delivered via a particle bombardment approach, many of which have been described.

In one illustrative example, gas-driven particle acceleration can be achieved with devices such as those manufactured by Powderject Pharmaceuticals PLC (Oxford, UK) and Powderject Vaccines Inc. (Madison, WI), some examples of which are described in U.S. Patent Nos. 5,846,796; 6,010,478; 5,865,796; 5,584,807; and EP Patent No. 0500  
5 799. This approach offers a needle-free delivery approach wherein a dry powder formulation of microscopic particles, such as polynucleotide or polypeptide particles, are accelerated to high speed within a helium gas jet generated by a hand held device, propelling the particles into a target tissue of interest.

In a related embodiment, other devices and methods that may be useful  
10 for gas-driven needle-less injection of compositions of the present invention include those provided by Bioject, Inc. (Portland, OR), some examples of which are described in U.S. Patent Nos. 4,790,824; 5,064,413; 5,312,335; 5,383,851; 5,399,163; 5,520,639 and 5,993,412.

According to another embodiment, the pharmaceutical compositions  
15 described herein will comprise one or more immunostimulants in addition to the immunogenic polynucleotide, polypeptide, antibody, T-cell and/or APC compositions of this invention. An immunostimulant refers to essentially any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an exogenous antigen. One preferred type of immunostimulant comprises an adjuvant. Many  
20 adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Certain adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant  
25 65 (Merck and Company, Inc., Rahway, NJ); AS-2 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as

GM-CSF, interleukin-2, -7, -12, and other like growth factors, may also be used as adjuvants.

Within certain embodiments of the invention, the adjuvant composition is preferably one that induces an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (e.g., IFN- $\gamma$ , TNF $\alpha$ , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (e.g., IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Certain preferred adjuvants for eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A, together with an aluminum salt. MPL® adjuvants are available from Corixa Corporation (Seattle, WA; see, for example, US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Patent Nos. 6,008,200 and 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato et al., *Science* 273:352, 1996. Another preferred adjuvant comprises a saponin, such as Quil A, or derivatives thereof, including QS21 and QS7 (Aquila Biopharmaceuticals Inc., Framingham, MA); Eacin; Digitonin; or *Gypsophila* or *Chenopodium quinoa* saponins. Other preferred formulations include more than one saponin in the adjuvant combinations of the present invention, for example combinations of at least two of the following group comprising QS21, QS7, Quil A,  $\beta$ -escin, or digitonin.

Alternatively the saponin formulations may be combined with vaccine vehicles composed of chitosan or other polycationic polymers, polylactide and polylactide-co-glycolide particles, poly-N-acetyl glucosamine-based polymer matrix, particles composed of polysaccharides or chemically modified polysaccharides, liposomes and lipid-based particles, particles composed of glycerol monoesters, etc. The saponins may also be formulated in the presence of cholesterol to form particulate structures such as liposomes or ISCOMs. Furthermore, the saponins may be formulated together with a polyoxyethylene ether or ester, in either a non-particulate solution or suspension, or in a particulate structure such as a paucilamellar liposome or ISCOM. The saponins may also be formulated with excipients such as Carbopol<sup>®</sup> to increase viscosity, or may be formulated in a dry powder form with a powder excipient such as lactose.

In one preferred embodiment, the adjuvant system includes the combination of a monophosphoryl lipid A and a saponin derivative, such as the combination of QS21 and 3D-MPL<sup>®</sup> adjuvant, as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and tocopherol. Another particularly preferred adjuvant formulation employing QS21, 3D-MPL<sup>®</sup> adjuvant and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Another enhanced adjuvant system involves the combination of a CpG-containing oligonucleotide and a saponin derivative particularly the combination of CpG and QS21 is disclosed in WO 00/09159. Preferably the formulation additionally comprises an oil in water emulsion and tocopherol.

Additional illustrative adjuvants for use in the pharmaceutical compositions of the invention include Montanide ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (e.g., SBAS-2 or SBAS-4, available from SmithKline Beecham, Rixensart, Belgium), Detox (Enhazyn<sup>®</sup>) (Corixa, Hamilton, MT), RC-529 (Corixa, Hamilton, MT) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described

in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosure of which are incorporated herein by reference in their entireties, and polyoxyethylene ether adjuvants such as those described in WO 99/52549A1.

Other preferred adjuvants include adjuvant molecules of the general  
5 formula



wherein,  $n$  is 1-50,  $A$  is a bond or  $-\text{C}(\text{O})-$ ,  $R$  is  $\text{C}_{1-50}$  alkyl or Phenyl  $\text{C}_{1-50}$  alkyl.

One embodiment of the present invention consists of a vaccine formulation comprising a polyoxyethylene ether of general formula (I), wherein  $n$  is  
10 between 1 and 50, preferably 4-24, most preferably 9; the  $R$  component is  $\text{C}_{1-50}$ , preferably  $\text{C}_4\text{-C}_{20}$  alkyl and most preferably  $\text{C}_{12}$  alkyl, and  $A$  is a bond. The concentration of the polyoxyethylene ethers should be in the range 0.1-20%, preferably from 0.1-10%, and most preferably in the range 0.1-1%. Preferred polyoxyethylene ethers are selected from the following group: polyoxyethylene-9-lauryl ether,  
15 polyoxyethylene-9-stearyl ether, polyoxyethylene-8-stearyl ether, polyoxyethylene-4-lauryl ether, polyoxyethylene-35-lauryl ether, and polyoxyethylene-23-lauryl ether. Polyoxyethylene ethers such as polyoxyethylene lauryl ether are described in the Merck index (12<sup>th</sup> edition: entry 7717). These adjuvant molecules are described in WO 99/52549.

20 The polyoxyethylene ether according to the general formula (I) above may, if desired, be combined with another adjuvant. For example, a preferred adjuvant combination is preferably with CpG as described in the pending UK patent application GB 9820956.2.

According to another embodiment of this invention, an immunogenic  
25 composition described herein is delivered to a host via antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be  
30 immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs

may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic  
5 cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (see Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*,  
10 with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As  
15 an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (see Zitvogel et al., *Nature Med.* 4:594-600, 1998).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph  
20 nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF $\alpha$  to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into  
25 dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF $\alpha$ , CD40 ligand, LPS, IL3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized  
30 phenotypes. However, this nomenclature should not be construed to exclude all

possible intermediate stages of differentiation. Immature dendritic cells are characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fcγ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (e.g., CD54 and CD11) and costimulatory molecules (e.g., CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide of the invention (or portion or other variant thereof) such that the encoded polypeptide, or an immunogenic portion thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a pharmaceutical composition comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and Cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the tumor polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (e.g., vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (e.g., a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will typically vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, mucosal, intravenous, intracranial, intraperitoneal, subcutaneous and intramuscular administration.



Carriers for use within such pharmaceutical compositions are biocompatible, and may also be biodegradable. In certain embodiments, the formulation preferably provides a relatively constant level of active component release. In other embodiments, however, a more rapid rate of release immediately upon administration may be desired. The formulation of such compositions is well within the level of ordinary skill in the art using known techniques. Illustrative carriers useful in this regard include microparticles of poly(lactide-co-glycolide), polyacrylate, latex, starch, cellulose, dextran and the like. Other illustrative delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (e.g., a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (see e.g., U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

In another illustrative embodiment, biodegradable microspheres (e.g., polylactate polyglycolate) are employed as carriers for the compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344, 5,407,609 and 5,942,252. Modified hepatitis B core protein carrier systems, such as described in WO/99 40934, and references cited therein, will also be useful for many applications. Another illustrative carrier/delivery system employs a carrier comprising particulate-protein complexes, such as those described in U.S. Patent No. 5,928,647, which are capable of inducing a class I-restricted cytotoxic T lymphocyte responses in a host.

The pharmaceutical compositions of the invention will often further comprise one or more buffers (e.g., neutral buffered saline or phosphate buffered saline), carbohydrates (e.g., glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (e.g., aluminum hydroxide), solutes that

render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate.

The pharmaceutical compositions described herein may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers are typically sealed in such a way to preserve the sterility and stability of the formulation until use. In general, formulations may be stored as suspensions, solutions or emulsions in oily or aqueous vehicles. Alternatively, a pharmaceutical composition may be stored in a freeze-dried condition requiring only the addition of a sterile liquid carrier immediately prior to use.

The development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including e.g., oral, parenteral, intravenous, intranasal, and intramuscular administration and formulation, is well known in the art, some of which are briefly discussed below for general purposes of illustration.

In certain applications, the pharmaceutical compositions disclosed herein may be delivered *via* oral administration to an animal. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet.

The active compounds may even be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like (see, for example, Mathiowitz *et al.*, *Nature* 1997 Mar 27;386(6623):410-4; Hwang *et al.*, *Crit Rev Ther Drug Carrier Syst* 1998;15(3):243-84; U. S. Patent 5,641,515; U. S. Patent 5,580,579 and U. S. Patent 5,792,451). Tablets, troches, pills, capsules and the like may also contain any of a variety of additional components, for example, a binder, such as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may

be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, 5 tablets, pills, or capsules may be coated with shellac, sugar, or both. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

Typically, these formulations will contain at least about 0.1% of the 10 active compound or more, although the percentage of the active ingredient(s) may, of course, be varied and may conveniently be between about 1 or 2% and about 60% or 70% or more of the weight or volume of the total formulation. Naturally, the amount of active compound(s) in each therapeutically useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. 15 Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

For oral administration the compositions of the present invention may 20 alternatively be incorporated with one or more excipients in the form of a mouthwash, dentifrice, buccal tablet, oral spray, or sublingual orally-administered formulation. Alternatively, the active ingredient may be incorporated into an oral solution such as one containing sodium borate, glycerin and potassium bicarbonate, or dispersed in a dentifrice, or added in a therapeutically-effective amount to a composition that may 25 include water, binders, abrasives, flavoring agents, foaming agents, and humectants. Alternatively the compositions may be fashioned into a tablet or solution form that may be placed under the tongue or otherwise dissolved in the mouth.

In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, intravenously, intramuscularly, or even 30 intraperitoneally. Such approaches are well known to the skilled artisan, some of which

are further described, for example, in U. S. Patent 5,543,158; U. S. Patent 5,641,515 and U. S. Patent 5,399,363. In certain embodiments, solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations generally will contain a preservative to prevent the growth of microorganisms.

Illustrative pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (for example, see U. S. Patent 5,466,468). In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and/or by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

In one embodiment, for parenteral administration in an aqueous solution, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will

be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. Moreover, for human administration, preparations will of course preferably meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

In another embodiment of the invention, the compositions disclosed herein may be formulated in a neutral or salt form. Illustrative pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective.

The carriers can further comprise any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human.

In certain embodiments, the pharmaceutical compositions may be delivered by intranasal sprays, inhalation, and/or other aerosol delivery vehicles.

Methods for delivering genes, nucleic acids, and peptide compositions directly to the lungs *via* nasal aerosol sprays has been described, *e.g.*, in U. S. Patent 5,756,353 and U. S. Patent 5,804,212. Likewise, the delivery of drugs using intranasal microparticle resins (Takenaga *et al.*, *J Controlled Release* 1998 Mar 2;52(1-2):81-7) and  
5 lysophosphatidyl-glycerol compounds (U. S. Patent 5,725,871) are also well-known in the pharmaceutical arts. Likewise, illustrative transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix is described in U. S. Patent 5,780,045.

In certain embodiments, liposomes, nanocapsules, microparticles, lipid particles, vesicles, and the like, are used for the introduction of the compositions of the present invention into suitable host cells/organisms. In particular, the compositions of  
10 the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like. Alternatively, compositions of the present invention can be bound, either covalently or non-covalently, to the surface of such carrier vehicles.

15 The formation and use of liposome and liposome-like preparations as potential drug carriers is generally known to those of skill in the art (see for example, Lasic, *Trends Biotechnol* 1998 Jul;16(7):307-21; Takakura, *Nippon Rinsho* 1998 Mar;56(3):691-5; Chandran *et al.*, *Indian J Exp Biol*. 1997 Aug;35(8):801-9; Margalit, *Crit Rev Ther Drug Carrier Syst*. 1995;12(2-3):233-61; U.S. Patent 5,567,434; U.S.  
20 Patent 5,552,157; U.S. Patent 5,565,213; U.S. Patent 5,738,868 and U.S. Patent 5,795,587, each specifically incorporated herein by reference in its entirety).

Liposomes have been used successfully with a number of cell types that are normally difficult to transfect by other procedures, including T cell suspensions, primary hepatocyte cultures and PC 12 cells (Renneisen *et al.*, *J Biol Chem*. 1990 Sep  
25 25;265(27):16337-42; Muller *et al.*, *DNA Cell Biol*. 1990 Apr;9(3):221-9). In addition, liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, various drugs, radiotherapeutic agents, enzymes, viruses, transcription factors, allosteric effectors and the like, into a variety of cultured cell lines and animals. Furthermore, the use of

liposomes does not appear to be associated with autoimmune responses or unacceptable toxicity after systemic delivery.

In certain embodiments, liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric  
5 bilayer vesicles (also termed multilamellar vesicles (MLVs)).

Alternatively, in other embodiments, the invention provides for pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds in a stable and reproducible way (see, for example, Quintanar-Guerrero *et al.*, Drug Dev Ind Pharm.  
10 1998 Dec;24(12):1113-28). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1  $\mu\text{m}$ ) may be designed using polymers able to be degraded *in vivo*. Such particles can be made as described, for example, by Couvreur *et al.*, Crit Rev Ther Drug Carrier Syst. 1988;5(1):1-20; zur Mühlen *et al.*, Eur J Pharm Biopharm. 1998 Mar;45(2):149-55; Zambaux *et al.* J  
15 Controlled Release. 1998 Jan 2;50(1-3):31-40; and U. S. Patent 5,145,684.

#### Cancer Therapeutic Methods

In further aspects of the present invention, the pharmaceutical compositions described herein may be used for the treatment of cancer, particularly for the immunotherapy of breast cancer. Within such methods, the pharmaceutical  
20 compositions described herein are administered to a patient, typically a warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Accordingly, the above pharmaceutical compositions may be used to prevent the development of a cancer or to treat a patient afflicted with a cancer. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical  
25 removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs. As discussed above, administration of the pharmaceutical compositions may be by any suitable method, including administration by intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal, intradermal, anal, vaginal, topical and oral routes.

Within certain embodiments, immunotherapy may be active immunotherapy, in which treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immune response-modifying agents (such as polypeptides and polynucleotides as provided  
5 herein).

Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host  
10 immune system. Examples of effector cells include T cells as discussed above, T lymphocytes (such as CD8<sup>+</sup> cytotoxic T lymphocytes and CD4<sup>+</sup> T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as dendritic cells and macrophages) expressing a polypeptide provided herein. T cell receptors and antibody  
15 receptors specific for the polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The polypeptides provided herein may also be used to generate antibodies or anti-idiotypic antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

Effector cells may generally be obtained in sufficient quantities for adoptive immunotherapy by growth *in vitro*, as described herein. Culture conditions for expanding single antigen-specific effector cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. Such *in vitro* culture conditions typically use intermittent stimulation with antigen, often in the presence of  
25 cytokines (such as IL-2) and non-dividing feeder cells. As noted above, immunoreactive polypeptides as provided herein may be used to rapidly expand antigen-specific T cell cultures in order to generate a sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast and/or B cells, may be pulsed with immunoreactive polypeptides  
30 or transfected with one or more polynucleotides using standard techniques well known



in the art. For example, antigen-presenting cells can be transfected with a polynucleotide having a promoter appropriate for increasing expression in a recombinant virus or other expression system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see, for example, Cheever et al., Immunological Reviews 157:177, 1997*).

Alternatively, a vector expressing a polypeptide recited herein may be introduced into antigen presenting cells taken from a patient and clonally propagated *in vivo* for transplant back into the same patient. Transfected cells may be reintroduced into the patient using any means known in the art, preferably in sterile form by intravenous, intracavitary, intraperitoneal or intratumor administration.

Routes and frequency of administration of the therapeutic compositions described herein, as well as dosage, will vary from individual to individual, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.*, untreated) level. Such response can be monitored by measuring the anti-tumor antibodies in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and vaccines comprising one or more polypeptides, the amount of each polypeptide present in a dose

ranges from about 2.5  $\mu$ g to 5 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a tumor protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

#### Cancer Detection and Diagnostic Compositions, Methods and Kits

In general, a cancer may be detected in a patient based on the presence of one or more breast tumor proteins and/or polynucleotides encoding such proteins in a biological sample (for example, blood, sera, sputum urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as breast cancer. In addition, such proteins may be useful for the detection of other cancers. The binding agents provided herein generally permit detection of the level of antigen that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding a tumor protein, which is also indicative of the presence or absence of a cancer. In general, a breast tumor sequence should be present at a level that is at least three fold higher in tumor tissue than in normal tissue.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, the presence or absence of a cancer in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b)

detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include full length breast tumor proteins and polypeptide portions thereof to which the binding agent binds, as described above.

The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is

preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 µg, and preferably about 100 ng to about 1 µg, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a detection reagent (preferably a second antibody capable of binding to a different site on the polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20<sup>TM</sup> (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as

phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of a cancer, such as breast cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average

mean signal obtained when the immobilized antibody is incubated with samples from patients without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (i.e., sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (i.e., the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for a cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the binding agent is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second, labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing the second binding agent flows through the membrane. The detection of bound second binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second binding agent and to the area of immobilized binding agent. Concentration of second binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized

on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 µg, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the tumor proteins or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use tumor polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such tumor protein specific antibodies may correlate with the presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with a tumor protein in a biological sample. Within certain methods, a biological sample comprising CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells isolated from a patient is incubated with a tumor polypeptide, a polynucleotide encoding such a polypeptide and/or an APC that expresses at least an immunogenic portion of such a polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with polypeptide (e.g., 5 - 25 µg/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of tumor polypeptide to serve as a control. For CD4<sup>+</sup> T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8<sup>+</sup> T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation

that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding a tumor protein in a biological sample. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of a tumor cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (i.e., hybridizes to) a polynucleotide encoding the tumor protein. The amplified cDNA is then separated and detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding a tumor protein may be used in a hybridization assay to detect the presence of polynucleotide encoding the tumor protein in a biological sample.

To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding a tumor protein of the invention that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above. Oligonucleotide primers and/or probes which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence as disclosed herein. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

One preferred assay employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological sample, such as biopsy tissue, and is reverse transcribed to produce cDNA molecules.



PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an individual who is not afflicted with a cancer. The amplification reaction may be  
5 performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered positive.

In another embodiment, the compositions described herein may be used  
10 as markers for the progression of cancer. In this embodiment, assays as described above for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) or polynucleotide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the  
15 level of polypeptide or polynucleotide detected increases over time. In contrast, the cancer is not progressing when the level of reactive polypeptide or polynucleotide either remains constant or decreases with time.

Certain *in vivo* diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound  
20 binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

As noted above, to improve sensitivity, multiple tumor protein markers may be assayed within a given sample. It will be apparent that binding agents specific  
25 for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be used concurrently. The selection of tumor protein markers may be based on routine experiments to determine combinations that results in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment. For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to a tumor protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding a tumor protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding a tumor protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a tumor protein.

The following Examples are offered by way of illustration and not by way of limitation.

#### EXAMPLE I

##### PREPARATION OF BREAST TUMOR-SPECIFIC CDNAS USING DIFFERENTIAL DISPLAY RT-PCR

This Example illustrates the preparation of cDNA molecules encoding breast tumor-specific polypeptides using a differential display screen.

#### A. Preparation of B18Ag1 cDNA and Characterization of mRNA Expression

Tissue samples were prepared from breast tumor and normal tissue of a patient with breast cancer that was confirmed by pathology after removal from the patient. Normal RNA and tumor RNA was extracted from the samples and mRNA was

isolated and converted into cDNA using a (dT)<sub>12</sub>AG (SEQ ID NO:130) anchored 3' primer. Differential display PCR was then executed using a randomly chosen primer (CTTCAACCTC) (SEQ ID NO:103). Amplification conditions were standard buffer containing 1.5 mM MgCl<sub>2</sub>, 20 pmol of primer, 500 pmol dNTP, and 1 unit of *Taq* DNA  
5 polymerase (Perkin-Elmer, Branchburg, NJ). Forty cycles of amplification were performed using 94°C denaturation for 30 seconds, 42°C annealing for 1 minute, and 72 °C extension for 30 seconds. An RNA fingerprint containing 76 amplified products was obtained. Although the RNA fingerprint of breast tumor tissue was over 98% identical to that of the normal breast tissue, a band was repeatedly observed to be  
10 specific to the RNA fingerprint pattern of the tumor. This band was cut out of a silver stained gel, subcloned into the T-vector (Novagen, Madison, WI) and sequenced.

The sequence of the cDNA, referred to as B18Ag1, is provided in SEQ ID NO:1. A database search of GENBANK and EMBL revealed that the B18Ag1 fragment initially cloned is 77% identical to the endogenous human retroviral element  
15 S71, which is a truncated retroviral element homologous to the Simian Sarcoma Virus (SSV). S71 contains an incomplete *gag* gene, a portion of the *pol* gene and an LTR-like structure at the 3' terminus (see Werner et al., *Virology* 174:225-238 (1990)). B18Ag1 is also 64% identical to SSV in the region corresponding to the P30 (*gag*) locus. B18Ag1 contains three separate and incomplete reading frames covering a region which  
20 shares considerable homology to a wide variety of *gag* proteins of retroviruses which infect mammals. In addition, the homology to S71 is not just within the *gag* gene, but spans several kb of sequence including an LTR.

B18Ag1-specific PCR primers were synthesized using computer analysis guidelines. RT-PCR amplification (94°C, 30 seconds; 60°C → 42°C, 30 seconds; 72°C,  
25 30 seconds for 40 cycles) confirmed that B18Ag1 represents an actual mRNA sequence present at relatively high levels in the patient's breast tumor tissue. The primers used in amplification were B18Ag1-1 (CTG CCT GAG CCA CAA ATG) (SEQ ID NO:128) and B18Ag1-4 (CCG GAG GAG GAA GCT AGA GGA ATA) (SEQ ID NO:129) at a 3.5 mM magnesium concentration and a pH of 8.5, and B18Ag1-2 (ATG GCT ATT  
30 TTC GGG GCC TGA CA) (SEQ ID NO:126) and B18Ag1-3 (CCG GTA TCT CCT

CGT GGG TAT T) (SEQ ID NO:127) at 2 mM magnesium at pH 9.5. The same experiments showed exceedingly low to nonexistent levels of expression in this patient's normal breast tissue (see Figure 1). RT-PCR experiments were then used to show that B18Ag1 mRNA is present in nine other breast tumor samples (from Brazilian and American patients) but absent in, or at exceedingly low levels in, the normal breast tissue corresponding to each cancer patient. RT-PCR analysis has also shown that the B18Ag1 transcript is not present in various normal tissues (including lymph node, myocardium and liver) and present at relatively low levels in PBMC and lung tissue. The presence of B18Ag1 mRNA in breast tumor samples, and its absence from normal breast tissue, has been confirmed by Northern blot analysis, as shown in Figure 2.

The differential expression of B18Ag1 in breast tumor tissue was also confirmed by RNase protection assays. Figure 3 shows the level of B18Ag1 mRNA in various tissue types as determined in four different RNase protection assays. Lanes 1-12 represent various normal breast tissue samples, lanes 13-25 represent various breast tumor samples; lanes 26-27 represent normal prostate samples; lanes 28-29 represent prostate tumor samples; lanes 30-32 represent colon tumor samples; lane 33 represents normal aorta; lane 34 represents normal small intestine; lane 35 represents normal skin, lane 36 represents normal lymph node; lane 37 represents normal ovary; lane 38 represents normal liver; lane 39 represents normal skeletal muscle; lane 40 represents a first normal stomach sample, lane 41 represents a second normal stomach sample; lane 42 represents a normal lung; lane 43 represents normal kidney; and lane 44 represents normal pancreas. Interexperimental comparison was facilitated by including a positive control RNA of known  $\beta$ -actin message abundance in each assay and normalizing the results of the different assays with respect to this positive control.

RT-PCR and Southern Blot analysis has shown the B18Ag1 locus to be present in human genomic DNA as a single copy endogenous retroviral element. A genomic clone of approximately 12-18 kb was isolated using the initial B18Ag1 sequence as a probe. Four additional subclones were also isolated by XbaI digestion. Additional retroviral sequences obtained from the ends of the XbaI digests of these clones (located as shown in Figure 4) are shown as SEQ ID NO:3 - SEQ ID NO:10.

where SEQ ID NO:3 shows the location of the sequence labeled 10 in Figure 4, SEQ ID NO:4 shows the location of the sequence labeled 11-29, SEQ ID NO:5 shows the location of the sequence labeled 3, SEQ ID NO:6 shows the location of the sequence labeled 6, SEQ ID NO:7 shows the location of the sequence labeled 12, SEQ ID NO:8 shows the location of the sequence labeled 13, SEQ ID NO:9 shows the location of the sequence labeled 14 and SEQ ID NO:10 shows the location of the sequence labeled 11-22.

Subsequent studies demonstrated that the 12-18 kb genomic clone contains a retroviral element of about 7.75 kb, as shown in Figures 5A and 5B. The sequence of this retroviral element is shown in SEQ ID NO:141. The numbered line at the top of Figure 5A represents the sense strand sequence of the retroviral genomic clone. The box below this line shows the position of selected restriction sites. The arrows depict the different overlapping clones used to sequence the retroviral element. The direction of the arrow shows whether the single-pass subclone sequence corresponded to the sense or anti-sense strand. Figure 5B is a schematic diagram of the retroviral element containing B18Ag1 depicting the organization of viral genes within the element. The open boxes correspond to predicted reading frames, starting with a methionine, found throughout the element. Each of the six likely reading frames is shown, as indicated to the left of the boxes, with frames 1-3 corresponding to those found on the sense strand.

Using the cDNA of SEQ ID NO:1 as a probe, a longer cDNA was obtained (SEQ ID NO:227) which contains minor nucleotide differences (less than 1%) compared to the genomic sequence shown in SEQ ID NO:141.

B. Preparation of cDNA Molecules Encoding Other Breast Tumor-Specific Polypeptides

Normal RNA and tumor RNA was prepared and mRNA was isolated and converted into cDNA using a (dT)<sub>12</sub>AG anchored 3' primer, as described above. Differential display PCR was then executed using the randomly chosen primers of SEQ ID NOs:87-125. Amplification conditions were as noted above, and bands observed to be specific to the RNA fingerprint pattern of the tumor were cut out of a silver stained

gel, subcloned into either the T-vector (Novagen, Madison, WI) or the pCRII vector (Invitrogen, San Diego, CA) and sequenced. The sequences are provided in SEQ ID NO:11 - SEQ ID NO:86. Of the 79 sequences isolated, 67 were found to be novel (SEQ ID NOs:11-26 and 28-77) (*see also* Figures 6-20).

- 5 An extended DNA sequence (SEQ ID NO:290) for the antigen B15Ag1 (originally identified partial sequence provided in SEQ ID NO:27) was obtained in further studies. Comparison of the sequence of SEQ ID NO:290 with those in the gene bank as described above, revealed homology to the known human  $\beta$ -A actinin gene. Further studies led to the isolation of the full-length cDNA sequence for the antigen
- 10 B21GT2 (also referred to as B311D; originally identified partial cDNA sequence provided in SEQ ID NOs:56). The full-length sequence is provided in SEQ ID NO:307, with the corresponding amino acid sequence being provided in SEQ ID NO:308. Further studies led to the isolation of a splice variant of B311D. The B311D clone of SEQ ID NO:316 was sequenced and a XhoI/NotI fragment from this clone was gel
- 15 purified and 32P-cdTP labeled by random priming for use as a probe for further screening to obtain additional B311D gene sequence. Two fractions of a human breast tumor cDNA bacterial library were screened using standard techniques. One of the clones isolated in this manner yielded additional sequence which includes a poly A+ tail. The determined cDNA sequence of this clone (referred to as B311D\_BT1\_1A) is
- 20 provided in SEQ ID NO:317. The sequences of SEQ ID NOs:316 and 317 were found to share identity over a 464 bp region, with the sequences diverging near the poly A+ sequence of SEQ ID NO:317.

- Subsequent studies identified an additional 146 sequences (SEQ ID NOs:142-289), of which 115 appeared to be novel (SEQ ID NOs:142, 143, 146-152,
- 25 154-166, 168-176, 178-192, 194-198, 200-204, 206, 207, 209-214, 216, 218, 219, 221-240, 243-245, 247, 250, 251, 253, 255, 257-266, 268, 269, 271-273, 275, 276, 278, 280, 281, 284, 288 and 291). To the best of the inventors' knowledge none of the previously identified sequences have heretofore been shown to be expressed at a greater level in human breast tumor tissue than in normal breast tissue.

In further studies, several different splice forms of the antigen B11Ag1 (also referred to as B305D) were isolated, with each of the various splice forms containing slightly different versions of the B11Ag1 coding frame. Splice junction sequences define individual exons which, in various patterns and arrangements, make up the various splice forms. Primers were designed to examine the expression pattern of each of the exons using RT-PCR as described below. Each exon was found to show the same expression pattern as the original B11Ag1 clone, with expression being breast tumor-, normal prostate- and normal testis-specific. The determined cDNA sequences for the isolated protein coding exons are provided in SEQ ID NOs:292-298, respectively. The predicted amino acid sequences corresponding to the sequences of SEQ ID NOs:292 and 298 are provided in SEQ ID NOs:299 and 300. Additional studies using rapid amplification of cDNA ends (RACE), a 5' specific primer to one of the splice forms of B11Ag1 provided above and a breast adenocarcinoma, led to the isolation of three additional, related, splice forms referred to as isoforms B11C-15, B11C-8 and B11C-9,16. The determined cDNA sequences for these isoforms are provided in SEQ ID NO: 301-303, with the corresponding predicted amino acid sequences being provided in SEQ ID NOs:304-306.

The protein coding region of B11C-15 (SEQ ID NO: 301; also referred to as B305D isoform C) was used as a query sequence in a BLASTN search of the Genbank DNA database. A match was found to a genomic clone from chromosome 21 (Accession no. AP001465). The pairwise alignments provided in the BLASTN output were used to identify the putative exon, or coding, sequence of the chromosome 21 sequence that corresponds to the B305D sequence. Based on the BlastN pairwise alignments, the following pieces of GenBank record AP001465 were put together: base pairs 67978-68499, 72870-72987, 73144-73335, 76085-76206, 77905-78085, 80520-80624, 87602-87633. This sequence was then aligned with the B305D isoform C sequence using the DNA Star Seqman program and excess sequence was deleted in such a way as to maintain the sequence most similar to B305D. The final edited form of the chromosome 21 sequence was 96.5% identical to B305D. This resulting edited sequence from chromosome 21 was then translated and found to contain no stop codons

other than the final stop codon in the same position as that for B305D. As with B305D, the chromosome 21 sequence (provided in SEQ ID NO: 325) encoded a protein (SEQ ID NO: 326) with 384 amino acids. An alignment of this protein with the B305D isoform C protein (SEQ ID NO: 304) showed 90% amino acid identity.

5 The cDNA sequence of B305D isoform C (SEQ ID NO: 301) was used to identify homologs by searching the High Throughput Genome Sequencing (HTGS) database (NCBI, National Institutes for Health, Bethesda, MD). Homologs were identified on Chromosome 2 (Clone ID 9838181), Chromosome 10 (Clone ID 10933022), Chromosome 15 (Clone ID 11560284). These homologs shared greater  
10 than 90% identity with B305D isoform C at the nucleic acid level. All three of these homologs encode 384 amino acid ORFs that share greater than 90% identity with the amino acid sequence of SEQ ID NO: 304. Further searching of the GenBank database with the sequence of SEQ ID NO: 301 yielded a partial sequence homolog on Chromosome 22 (Clone ID 5931507). cDNA sequences for the Chromosome 2, 10, 15  
15 and 22 homologs were constructed based on the homology with B305D isoform C and the conserved sequences at intron-exon junctions. The cDNA sequences for the Chromosome 22, 2, 15 and 10 homologs are provided in SEQ ID NO: 327-330, respectively, with the corresponding amino acid sequences being provided in SEQ ID NO: 331, 334, 333 and 332, respectively.

20 In subsequent studies on B305D isoform A (cDNA sequence provided in SEQ ID NO:292), the cDNA sequence (provided in SEQ ID NO:313) was found to contain an additional guanine residue at position 884, leading to a frameshift in the open reading frame. The determined DNA sequence of this ORF is provided in SEQ ID NO:314. This frameshift generates a protein sequence (provided in SEQ ID NO:315) of  
25 293 amino acids that contains the C-terminal domain common to the other isoforms of B305D but that differs in the N-terminal region.



## EXAMPLE 2

## PREPARATION OF B18Ag1 DNA FROM HUMAN GENOMIC DNA

This Example illustrates the preparation of B18Ag1 DNA by  
5 amplification from human genomic DNA.

B18Ag1 DNA may be prepared from 250 ng human genomic DNA using  
20 pmol of B18Ag1 specific primers, 500 pmol dNTPS and 1 unit of *Taq* DNA  
polymerase (Perkin Elmer, Branchburg, NJ) using the following amplification  
parameters: 94°C for 30 seconds denaturing, 30 seconds 60°C to 42°C touchdown  
10 annealing in 2°C increments every two cycles and 72°C extension for 30 seconds. The  
last increment (a 42°C annealing temperature) should cycle 25 times. Primers were  
selected using computer analysis. Primers synthesized were B18Ag1-1, B18Ag1-2,  
B18Ag1-3, and B18Ag1-4. Primer pairs that may be used are 1+3, 1+4, 2+3, and 2+4.

Following gel electrophoresis, the band corresponding to B18Ag1 DNA  
15 may be excised and cloned into a suitable vector.

## EXAMPLE 3

## PREPARATION OF B18Ag1 DNA FROM BREAST TUMOR cDNA

This Example illustrates the preparation of B18Ag1 DNA by  
20 amplification from human breast tumor cDNA.

First strand cDNA is synthesized from RNA prepared from human breast  
tumor tissue in a reaction mixture containing 500 ng poly A<sup>+</sup> RNA, 200 pmol of the  
primer (I)<sub>12</sub>AG (i.e., TTT TTT TTT TTT AG) (SEQ ID NO:130), 1X first strand  
25 reverse transcriptase buffer, 6.7 mM DTT, 500 nmol dNTPs, and 1 unit AMV or  
MMLV reverse transcriptase (from any supplier, such as Gibco-BRL (Grand Island,  
NY)) in a final volume of 30 µL. After first strand synthesis, the cDNA is diluted  
approximately 25 fold and 1 µL is used for amplification as described in Example 2.  
While some primer pairs can result in a heterogeneous population of transcripts, the  
30 primers B18Ag1-2 (5'-ATG GCT ATT TTC GGG GGC TGA CA) (SEQ ID NO:126)

and B18Ag1-3 (5'CCG GTA TCT CCT CGT GGG TAT T) (SEQ ID NO:127) yield a single 151 bp amplification product.

#### EXAMPLE 4

##### 5 IDENTIFICATION OF B-CELL AND T-CELL EPITOPES OF B18Ag1

This Example illustrates the identification of B18Ag1 epitopes.

The B18Ag1 sequence can be screened using a variety of computer algorithms. To determine B-cell epitopes, the sequence can be screened for  
10 hydrophobicity and hydrophilicity values using the method of Hopp, *Prog. Clin. Biol. Res.* 172B:367-77 (1985) or, alternatively, Cense et al., *J. Exp. Med.* 164:1779-84 (1986) or Sponge et al., *J. Immunol.* 138:204-12 (1987). Additional Class II MHC (antibody or B-cell) epitopes can be predicted using programs such as AMPHI (e.g., Margalit et al., *J. Immunol.* 138:2213 (1987)) or the methods of Rothbard and Taylor  
15 (e.g., *EMBO J.* 7:93 (1988)).

Once peptides (15-20 amino acids long) are identified using these techniques, individual peptides can be synthesized using automated peptide synthesis equipment (available from manufacturers such as Perkin Elmer/Applied Biosystems Division, Foster City, CA) and techniques such as Merrifield synthesis. Following  
20 synthesis, the peptides can used to screen sera harvested from either normal or breast cancer patients to determine whether patients with breast cancer possess antibodies reactive with the peptides. Presence of such antibodies in breast cancer patient would confirm the immunogenicity of the specific B-cell epitope in question. The peptides can also be tested for their ability to generate a serologic or humoral immune in animals  
25 (mice, rats, rabbits, chimps etc.) following immunization *in vivo*. Generation of a peptide-specific antiserum following such immunization further confirms the immunogenicity of the specific B-cell epitope in question.

To identify T-cell epitopes, the B18Ag1 sequence can be screened using different computer algorithms which are useful in identifying 8-10 amino acid motifs  
30 within the B18Ag1 sequence which are capable of binding to HLA Class I MHC

molecules. (see, e.g., Rammensee et al., *Immunogenetics* 41:178-228 (1995)). Following synthesis such peptides can be tested for their ability to bind to class I MHC using standard binding assays (e.g., Sette et al., *J. Immunol.* 153:5586-92 (1994)) and more importantly can be tested for their ability to generate antigen reactive cytotoxic T-

5 cells following *in vitro* stimulation of patient or normal peripheral mononuclear cells using, for example, the methods of Bakker et al., *Cancer Res.* 55:5330-34 (1995); Visseren et al., *J. Immunol.* 154:3991-98 (1995); Kawakami et al., *J. Immunol.* 154:3961-68 (1995); and Kast et al., *J. Immunol.* 152:3904-12 (1994). Successful

10 *in vitro* generation of T-cells capable of killing autologous (bearing the same Class I MHC molecules) tumor cells following *in vitro* peptide stimulation further confirms the immunogenicity of the B18Ag1 antigen. Furthermore, such peptides may be used to generate murine peptide and B18Ag1 reactive cytotoxic T-cells following *in vivo* immunization in mice rendered transgenic for expression of a particular human MHC Class I haplotype (Vitello et al., *J. Exp. Med.* 173:1007-15 (1991)).

15 A representative list of predicted B18Ag1 B-cell and T-cell epitopes, broken down according to predicted HLA Class I MHC binding antigen, is shown below:

Predicted Th Motifs (B-cell epitopes) (SEQ ID NOS.: 131-133)

SSGRTFDDFHRYLLVGI

20 QGAAQKPINLSKXIEVVQGHDE

SPGVFLEHLQEAYRIYTPDLSA

Predicted HLA A2.1 Motifs (T-cell epitopes) (SEQ ID NOS.: 134-140)

YLLVGIQGA

25 GAAQKPINL

NLSKXIEVV

EVVQGHDES

HLQEAYRIY

NLAFAVAQAA

30 FVAQAAPDS

## EXAMPLE 5

## IDENTIFICATION OF T-CELL EPITOPES OF B11Ag1

This Example illustrates the identification of B11Ag1 (also referred to as B305D) epitopes. Four peptides, referred to as B11-8, B11-1, B11-5 and B11-12 (SEQ  
5 ID NOs:309-312, respectfully) were derived from the B11Ag1 gene.

Human CD8 T cells were primed *in vitro* to the peptide B11-8 using dendritic cells according to the protocol of Van Tsai et al. (*Critical Reviews in Immunology* 18:65-75, 1998). The resulting CD8 T cell cultures were tested for their ability to recognize the B11-8 peptide or a negative control peptide, presented by the B-  
10 LCL line, JY. Briefly, T cells were incubated with autologous monocytes in the presence of 10 ug/ml peptide, 10 ng/ml IL-7 and 10 ug/ml IL-2, and assayed for their ability to specifically lyse target cells in a standard 51-Cr release assay. As shown in Fig. 22, the bulk culture line demonstrated strong recognition of the B11-8 peptide with weaker recognition of the peptide B11-1.

15 A clone from this CTL line was isolated following rapid expansion using the monoclonal antibody OKT3 and human IL-2. As shown in Fig. 23, this clone (referred to as A1), in addition to being able to recognize specific peptide, recognized JY LCL transduced with the B11Ag1 gene. This data demonstrates that B11-8 is a naturally processed epitope of the B11Ag1 gene. In addition these T cells were further  
20 found to recognize and lyse, in an HLA-A2 restricted manner, an established tumor cell line naturally expressing B11Ag1 (Fig. 24). The T cells strongly recognize a lung adenocarcinoma (LT-140-22) naturally expressing B11Ag1 transduced with HLA-A2, as well as an A2+ breast carcinoma (CAMA-1) transduced with B11Ag1, but not untransduced lines or another negative tumor line (SW620).

25 These data clearly demonstrate that these human T cells recognize not only B11-specific peptides but also transduced cells, as well as naturally expressing tumor lines.

CTL lines raised against the antigens B11-5 and B11-12, using the procedures described above, were found to recognize corresponding peptide-coated  
30 targets.

## EXAMPLE 6

CHARACTERIZATION OF BREAST TUMOR GENES DISCOVERED BY  
DIFFERENTIAL DISPLAY PCR

5           The specificity and sensitivity of the breast tumor genes discovered by differential display PCR were determined using RT-PCR. This procedure enabled the rapid evaluation of breast tumor gene mRNA expression semiquantitatively without using large amounts of RNA. Using gene specific primers, mRNA expression levels in a variety of tissues were examined, including 8 breast tumors, 5 normal breasts, 2  
10 prostate tumors, 2 colon tumors, 1 lung tumor, and 14 other normal adult human tissues, including normal prostate, colon, kidney, liver, lung, ovary, pancreas, skeletal muscle, skin, stomach and testes.

To ensure the semiquantitative nature of the RT-PCR,  $\beta$ -actin was used as internal control for each of the tissues examined. Serial dilutions of the first strand  
15 cDNAs were prepared and RT-PCR assays performed using  $\beta$ -actin specific primers. A dilution was then selected that enabled the linear range amplification of  $\beta$ -actin template, and which was sensitive enough to reflect the difference in the initial copy number. Using this condition, the  $\beta$ -actin levels were determined for each reverse transcription reaction from each tissue. DNA contamination was minimized by DNase  
20 treatment and by assuring a negative result when using first strand cDNA that was prepared without adding reverse transcriptase.

Using gene specific primers, the mRNA expression levels were determined in a variety of tissues. To date, 38 genes have been successfully examined by RT-PCR, five of which exhibit good specificity and sensitivity for breast tumors  
25 (B15AG-1, B31GA1b, B38GA2a, B11A1a and B18AG1a). Figures 21A and 21B depict the results for three of these genes: B15AG-1 (SEQ ID NO:27), B31GA1b (SEQ ID NO:148) and B38GA2a (SEQ ID NO:157). Table I summarizes the expression level of all the genes tested in normal breast tissue and breast tumors, and also in other tissues.

TABLE I

Percentage of Breast Cancer Antigens that are Expressed in Various Tissues

5	Breast Tissues	Over-expressed in Breast Tumors	84%
		Equally Expressed in Normals and Tumor	16%
10	Other Tissues	Over-expressed in Breast Tumors but not in any Normal Tissues	9%
		Over-expressed in Breast Tumors but Expressed in Some Normal Tissues	30%
15		Over-expressed in Breast Tumors but Equally Expressed in All Other Tissues	61%

20

## EXAMPLE 7

## PREPARATION AND CHARACTERIZATION OF ANTIBODIES AGAINST BREAST TUMOR

## POLYPEPTIDES

25 Polyclonal antibodies against the breast tumor antigen B305D were prepared as follows.

The breast tumor antigen expressed in an *E. coli* recombinant expression system was grown overnight in LB broth with the appropriate antibiotics at 37 °C in a shaking incubator. The next morning, 10 ml of the overnight culture was added to 500 ml to 2x YT plus appropriate antibiotics in a 2L-baffled Erlenmeyer flask. When the  
 30 Optical Density (at 560 nm) of the culture reached 0.4-0.6, the cells were induced with IPTG (1 mM). Four hours after induction with IPTG, the cells were harvested by centrifugation. The cells were then washed with phosphate buffered saline and centrifuged again. The supernatant was discarded and the cells were either frozen for future use or immediately processed. Twenty ml of lysis buffer was added to the cell  
 35 pellets and vortexed. To break open the *E. coli* cells, this mixture was then run through the French Press at a pressure of 16,000 psi. The cells were then centrifuged again and

the supernatant and pellet were checked by SDS-PAGE for the partitioning of the recombinant protein. For proteins that localized to the cell pellet, the pellet was resuspended in 10 mM Tris pH 8.0, 1% CHAPS and the inclusion body pellet was washed and centrifuged again. This procedure was repeated twice more. The washed  
5 inclusion body pellet was solubilized with either 8 M urea or 6 M guanidine HCl containing 10 mM Tris pH 8.0 plus 10 mM imidazole. The solubilized protein was added to 5 ml of nickel-chelate resin (Qiagen) and incubated for 45 min to 1 hour at room temperature with continuous agitation. After incubation, the resin and protein mixture were poured through a disposable column and the flow through was collected.  
10 The column was then washed with 10-20 column volumes of the solubilization buffer. The antigen was then eluted from the column using 8M urea, 10 mM Tris pH 8.0 and 300 mM imidazole and collected in 3 ml fractions. A SDS-PAGE gel was run to determine which fractions to pool for further purification.

As a final purification step, a strong anion exchange resin such as  
15 HiPrepQ (Biorad) was equilibrated with the appropriate buffer and the pooled fractions from above were loaded onto the column. Antigen was eluted off the column with a increasing salt gradient. Fractions were collected as the column was run and another SDS-PAGE gel was run to determine which fractions from the column to pool. The pooled fractions were dialyzed against 10 mM Tris pH 8.0. The protein was then vialled  
20 after filtration through a 0.22 micron filter and the antigens were frozen until needed for immunization.

Four hundred micrograms of B305D antigen was combined with 100 micrograms of muramyl dipeptide (MDP). Every four weeks rabbits were boosted with 100 micrograms mixed with an equal volume of Incomplete Freund's Adjuvant (IFA).  
25 Seven days following each boost, the animal was bled. Sera was generated by incubating the blood at 4 °C for 12-24 hours followed by centrifugation.

Ninety-six well plates were coated with B305D antigen by incubating with 50 microliters (typically 1 microgram) of recombinant protein at 4 °C for 20 hours. 250 microliters of BSA blocking buffer was added to the wells and incubated at room  
30 temperature for 2 hours. Plates were washed 6 times with PBS/0.01% Tween. Rabbit

sera was diluted in PBS. Fifty microliters of diluted sera was added to each well and incubated at room temperature for 30 min. Plates were washed as described above before 50 microliters of goat anti-rabbit horse radish peroxidase (HRP) at a 1:10000 dilution was added and incubated at room temperature for 30 min. Plates were again washed as described above and 100 microliters of TMB microwell peroxidase substrate was added to each well. Following a 15 min incubation in the dark at room temperature, the colorimetric reaction was stopped with 100 microliters of 1N H<sub>2</sub>SO<sub>4</sub> and read immediately at 450 nm. The polyclonal antibodies showed immunoreactivity to B305D.

Immunohistochemical (IHC) analysis of B305D expression in breast cancer and normal breast specimens was performed as follows. Paraffin-embedded formal fixed tissue was sliced into 8 micron sections. Steam heat induced epitope retrieval (SHIER) in 0.1 M sodium citrate buffer (pH 6.0) was used for optimal staining conditions. Sections were incubated with 10% serum/PBS for 5 minutes. Primary antibody was added to each section for 25 min at indicated concentrations followed by a 25 min incubation with either an anti-rabbit or anti-mouse biotinylated antibody. Endogenous peroxidase activity was blocked by three 1.5 min incubations with hydrogen peroxide. The avidin biotin complex/horseradish peroxidase (ABC/HRP) systems was used along with DAB chromagen to visualize antigen expression. Slides were counterstained with hematoxylin. B305D expression was detected in both breast tumor and normal breast tissue. However, the intensity of staining was much less in normal samples than in tumor samples and surface expression of B305D was observed only in breast tumor tissues.

A summary of real-time PCR and immunohistochemical analysis of B305D expression in an extensive panel of normal tissues is presented in Table II below. These results demonstrate minimal expression of B305D in testis, inconclusive results in gall bladder, and no detection in all other tissues tested.



TABLE II

miRNA	IHC staining	Tissue type	Summary
Moderately positive	Positive	Testis	Nuclear staining of small minority of spermatids; spermatozoa negative; seminoma negative
Negative	Negative	Thymus	No expression
N/A	Negative	Artery	No expression
Negative	Negative	Skeletal muscle	No expression
Negative	Positive (weak staining)	Small bowel	No expression
Negative	Positive (weak staining)	Ovary	No expression
Negative		Pituitary	No expression
Negative	Positive (weak staining)	Stomach	No expression
Negative	Negative	Spinal cord	No expression
Negative	Negative	Spleen	No expression
Negative	Negative	Ureter	No expression
N/A	Negative	Gall bladder	Inconclusive
N/A	Negative	Placenta	No expression
Negative	Negative	Thyroid	No expression
Negative	Negative	Heart	No expression
Negative	Negative	Kidney	No expression
Negative	Negative	Liver	No expression
Negative	Negative	Brain-cerebellum	No expression
Negative	Negative	Colon	No expression
Negative	Negative	Skin	No expression
Negative	Negative	Bone marrow	No expression
N/A	Negative	Parathyroid	No expression
Negative	Negative	Lung	No expression
Negative	Negative	Esophagus	No expression
Negative	Positive (weak staining)	Uterus	No expression
Negative	Negative	Adrenal	No expression
Negative	Negative	Pancreas	No expression
N/A	Negative	Lymph node	No expression
Negative	Negative	Brain-cortex	No expression
N/A	Negative	Fallopian tube	No expression
Negative	Positive (weak staining)	Bladder	No expression
Negative	N/A	Bone	No expression
Negative	N/A	Salivary gland	No expression

Negative	N/A	Activated PBMC	No expression
Negative	N/A	Resting PBMC	No expression
Negative	N/A	Trachea	No expression
Negative	N/A	Vena cava	No expression
Negative	N/A	Retina	No expression
Negative	N/A	Cartilage	No expression

## EXAMPLE 8

## PROTEIN EXPRESSION OF BREAST TUMOR ANTIGENS

5 This example describes the expression and purification of the breast tumor antigen B305D in *E. coli* and in mammalian cells.

Expression of B305D isoform C-15 (SEQ ID NO:301; translated to 384 amino acids) in *E. coli* was achieved by cloning the open reading frame of B305D isoform C-15 downstream of the first 30 amino acids of the *M. tuberculosis* antigen  
 10 Ra12 (SEQ ID NO:318) in pET17b. First, the internal EcoRI site in the B305D ORF was mutated without changing the protein sequence so that the gene could be cloned at the EcoRI site with Ra12. The PCR primers used for site-directed mutagenesis are shown in SEQ ID NO:319 (referred to as AW012) and SEQ ID NO:320 (referred to as AW013). The ORF of EcoRI site-modified B305D was then amplified by PCR using  
 15 the primers AW014 (SEQ ID NO:321) and AW015 (SEQ ID NO:322). The PCR product was digested with EcoRI and ligated to the Ra12/pET17b vector at the EcoRI site. The sequence of the resulting fusion construct (referred to as Ra12mB11C) was confirmed by DNA sequencing. The determined cDNA sequence for the fusion construct is provided in SEQ ID NO:323, with the amino acid sequence being provided  
 20 in SEQ ID NO:324.

The fusion construct was transformed into BL21(DE3)CodonPlus-RIL *E. coli* (Stratagene) and grown overnight in LB broth with kanamycin. The resulting culture was induced with IPTG. Protein was transferred to PVDF membrane and blocked with 5% non-fat milk (in PBS-Tween buffer), washed three times and  
 25 incubated with mouse anti-His tag antibody (Clontech) for 1 hour. The membrane was washed 3 times and probed with HRP-Protein A (Zymed) for 30 min. Finally, the

membrane was washed 3 times and developed with ECL (Amersham). Expression was detected by Western blot.

For recombinant expression in mammalian cells, B305D isoform C-15 (SEQ ID NO:301; translated to 384 amino acids) was subcloned into the mammalian expression vectors pCEP4 and pcDNA3.1 (Invitrogen). These constructs were transfected into HEK293 cells (ATCC) using Fugene 6 reagent (Roche). Briefly, the HEK cells were plated at a density of 100,000 cells/ml in DMEM (Gibco) containing 10% FBS (Hyclone) and grown overnight. The following day, 2  $\mu$ l of Fugene 6 was added to 100  $\mu$ l of DMEM containing no FBS and incubated for 15 minutes at room temperature. The Fugene 6/DMEM mixture was added to 1  $\mu$ g of B305D/pCEP4 or B305D/pcDNA plasmid DNA and incubated for 15 minutes at room temperature. The Fugene/DNA mix was then added to the HEK293 cells and incubated for 48-72 hours at 37 °C with 7% CO<sub>2</sub>. Cells were rinsed with PBS, the collected and pelleted by centrifugation.

For Western blot analysis, whole cell lysates were generated by incubating the cells in Triton-X100 containing lysis buffer for 30 minutes on ice. Lysates were then cleared by centrifugation at 10,000 rpm for 5 minutes at 4 °C. Samples were diluted with SDS\_PAGE loading buffer containing beta-mercaptoethanol, and boiled for 10 minutes prior to loading the SDS\_PAGE gel. Proteins were transferred to nitrocellulose and probed using Protein A purified anti-B305D rabbit polyclonal sera (prepared as described above) at a concentration of 1  $\mu$ g/ml. The blot was revealed with a goat anti-rabbit Ig coupled to HRP followed by incubation in ECL substrate. Expression of B305D was detected in the the HEK293 lysates transfected with B305D, but not in control HEK293 cells transfected with vector alone.

For FACS analysis, cells were washed further with ice cold staining buffer and then incubated with a 1:100 dilution of a goat anti-rabbit Ig (H+L)-FITC reagent (Southern Biotechnology) for 30 minutes on ice. Following 3 washes, the cells were resuspended in staining buffer containing Propidium Iodide (PI), a vital stain that

allows for identification of permeable cells, and then analyzed by FACS. The FACS analysis showed surface expression of B305D protein.

From the foregoing it will be appreciated that, although specific  
5 embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

## CLAIMS

## What is Claimed:

1. An isolated polynucleotide comprising a sequence selected from the group consisting of:

(a) sequences provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(b) complements of the sequences provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(d) sequences that hybridize to a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330, under moderately stringent conditions;

(e) sequences having at least 75% identity to a sequence of SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330;

(f) sequences having at least 90% identity to a sequence of SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330; and

(g) degenerate variants of a sequence provided in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330.

2. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) SEQ ID NO: 299, 300, 304-306, 308-312, 314, 326 and 331-334;

(b) sequences encoded by a polynucleotide of claim 1;

(c) sequences having at least 70% identity to a sequence encoded by a polynucleotide of claim 1; and

(d) sequences having at least 90% identity to a sequence encoded by a polynucleotide of claim 1.

3. An expression vector comprising a polynucleotide of claim 1 operably linked to an expression control sequence.

4. A host cell transformed or transfected with an expression vector according to claim 3.

5. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a polypeptide of claim 2.

6. A method for detecting the presence of a cancer in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with a binding agent that binds to a polypeptide of claim 2;
- (c) detecting in the sample an amount of polypeptide that binds to the binding agent; and
- (d) comparing the amount of polypeptide to a predetermined cut-off value and therefrom determining the presence of a cancer in the patient.

7. A fusion protein comprising at least one polypeptide according to claim 2.

8. An oligonucleotide that hybridizes to a sequence recited in SEQ ID NO: 1, 3-86, 142-298, 301-303, 307, 313, 314, 316, 317, 325 and 327-330 under moderately stringent conditions.

9. A method for stimulating and/or expanding T cells specific for a tumor protein, comprising contacting T cells with at least one component selected from the group consisting of:

- (a) polypeptides according to claim 2;
- (b) polynucleotides according to claim 1; and
- (c) antigen-presenting cells that express a polypeptide according to

claim 2,

under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

10. An isolated T cell population, comprising T cells prepared according to the method of claim 9.

11. A composition comprising a first component selected from the group consisting of physiologically acceptable carriers and immunostimulants, and a second component selected from the group consisting of:

- (a) polypeptides according to claim 2;
- (b) polynucleotides according to claim 1;
- (c) antibodies according to claim 5;
- (d) fusion proteins according to claim 7;
- (e) T cell populations according to claim 10; and
- (f) antigen presenting cells that express a polypeptide according to

claim 2.

12. A method for stimulating an immune response in a patient, comprising administering to the patient a composition of claim 11.

13. A method for the treatment of a cancer in a patient, comprising administering to the patient a composition of claim 11.

14. A method for determining the presence of a cancer in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide according to claim 8;
- (c) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and
- (d) compare the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence of the cancer in the patient.


15. A diagnostic kit comprising at least one oligonucleotide according to claim 8.

16. A diagnostic kit comprising at least one antibody according to claim 5 and a detection reagent, wherein the detection reagent comprises a reporter group.

17. A method for inhibiting the development of a cancer in a patient, comprising the steps of:


- (a) incubating CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells isolated from a patient with at least one component selected from the group consisting of: (i) polypeptides according to claim 2; (ii) polynucleotides according to claim 1; and (iii) antigen presenting cells that express a polypeptide of claim 2, such that T cell proliferate;
  - (b) administering to the patient an effective amount of the proliferated T cells,
- and thereby inhibiting the development of a cancer in the patient.






*Fig. 1*


2/19



*Fig. 2*



*Fig. 3*

GENOMIC CLONE MAP*Fig. 4*